Dietary Exposure and Risk Assessment

DP#: 351961 and 352204



# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OPP OFFICIAL RECORD **HEALTH EFFECTS DIVISION** SCIENTIFIC DATA REVIEWS **EPA SERIES 361** 

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

## **MEMORANDUM**

DATE:

1-MAY-2008

**SUBJECT:** 

Difenoconazole: Section 18 Emergency Exemption for use in/on Almond

and Cucurbit Vegetables (Group 9); Acute and Chronic Dietary Exposure

and Risk Assessment

PC Code:

128847

**DP Barcode:** 

351961 and 352204

**Decision Nos:** 

392318 and 391843

**Registration Nos:** 

08CA12 and 08GA01

**Petition No.:** 

NA

**Regulatory Action:** 

Section 18

Risk Assessment

Single Chemical

Case No.:

NA

Type: TXR No.: Dietary NA

CAS No.:

119446-68-3

MRID No.:

NA

**40 CFR** 

§180.475

FROM:

Mohsen Sahafeyan, Chemist

Registration Action Branch 1/Health Effects Division (RAB1/HED): 7509P

THROUGH:

Thurston Morton, Chemist Ware Solling for Transfor Morton

Yvonne Barnes, Chemist Affart

Dietary Exposure Science Advisory Council (DESAC)

Health Effects Division; 7509P

And

George F. Kramer, Ph.D., Senior Chemist

**RAB1/HED**; 7509P

TO:

Mohsen Sahafeyan, Chemist

RAB1/HED; 7509P

And

William D. Wassell, Chemist

RAB1/HED; 7509P

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

## **Executive Summary**

Aggregate (food + water) acute and chronic dietary exposure and risk assessments were conducted using the Dietary Exposure Evaluation Model - Food Consumption Intake Database (DEEM-FCID™, ver. 2.03) model. This model uses food consumption data from the United States Department of Agriculture's (USDA's) Continuing Surveys of Food Intakes by Individuals (CSFII; 1994-1996 and 1998). The analyses were performed to support two Section 18 requests for the use of fungicide difenoconazole [1-[2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-ylmethyl]-1*H*-1,2,4-triazole] in/on almond nutmeat, almond hulls, and cucurbit vegetables (group 9) with recommended time-limited tolerances at 0.05 ppm, 5.0 ppm, and 1.0 ppm, respectively.

The unrefined (Tier 1) acute and chronic analyses assumed tolerance-level residues, 100% crop treated (CT), and empirical and DEEM<sup>TM</sup> (ver. 7.81) default processing factors for most commodities. The drinking water values used in the dietary risk assessment were provided by the Environmental Fate and Effects Division (EFED; Memo, I. Maher, 19-JUN-2007; DP# 333319). The resulting acute and chronic aggregate exposure estimates were not of concern to HED.

## Acute Dietary (food + water) Exposure and Risk Assessment

The resulting acute food and water exposure estimates were not of concern to HED (<100% of the acute population-adjusted dose (aPAD)) at the 95<sup>th</sup> percentile of the exposure distribution for U.S. general population (2.9% aPAD) and all population subgroups; the most highly exposed population subgroup was all-infants <1 year old with 9.0% aPAD.

#### Chronic Dietary (food + water) Exposure and Risk Assessment

The resulting chronic food and water exposure estimates were not of concern to HED (<100% of the chronic population-adjusted dose (cPAD)) for U.S. general population (23% cPAD) and all population sub-groups; the most highly exposed population subgroup was children 1-2 years old with 65% cPAD.

#### Cancer Dietary Exposure and Risk Assessment

A cancer dietary assessment was <u>not</u> conducted for difenoconazole because the cancer noobservable-adverse-effect-level (NOAEL) is higher than the chronic reference dose (RfD); therefore, the chronic dietary risk estimate is protective of any cancer effects.

## Triazoles Dietary Exposure and Risk Assessments

The aggregate dietary (food + water) acute and chronic dietary exposure analyses for difenoconazole metabolites 1,2,4- triazole (1,2,4-T) and triazole alanine (TA) and triazole acetic acid (TAA) from all registered and proposed triazole-based fungicides uses already include exposure to almonds at higher-level exposures by other triazole-based fungicides and are not of concern (Memo, M. Doherty, DP#322238, 1-NOV-2005). In the case of cucurbit crop group,

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

the difenoconazole residue of 1.0 ppm translates to higher exposures to triazole metabolites than from other traizole-based fungicides uses on cucurbit that are already covered in the last dietary assessment for triazoles. Although, the aggregate dietary risk assessment for triazoles are being updated to include this level of use on cucurbit, it can be said with a fairly high degree of certainty that adding these Section 18 uses will not result in any risk issues for the triazole metabolites from HED's perspective (email from M. Doherty, 5/2/08).

#### I. Introduction

Dietary risk assessment incorporates both exposure and toxicity of a given pesticide. For acute and chronic assessments, the risk is expressed as a percentage of a maximum acceptable dose (i.e., the dose which HED has concluded will result in no unreasonable adverse health effects). This dose is referred to as the PAD. The PAD is equivalent to the point of departure (POD, NOAEL, LOAEL, e.g.) divided by the required uncertainty or safety factors.

For acute and non-cancer chronic exposures, HED is concerned when estimated dietary risk exceeds 100% of the PAD. HED is generally concerned when estimated cancer risk exceeds one in one million. References which discuss the acute and chronic risk assessments in more detail are available on the EPA/pesticides web site: "Available Information on Assessing Exposure from Pesticides, A User's Guide," 21-JUN-2000, web link: <a href="http://www.epa.gov/fedrgstr/EPA-PEST/2000/July/Day-12/6061.pdf">http://www.epa.gov/fedrgstr/EPA-PEST/2000/July/Day-12/6061.pdf</a>; or see SOP 99.6 (20-AUG-1999).

The most recent dietary risk assessment for difenoconazole was conducted by M. Sahafeyan (memo, D341303, 10/30/07).

#### II. Residue Information

Difenoconazole tolerances are published in 40 CFR§180.475 (Note: The correct sugar beet tolerance is 0.3 ppm instead of 0.01 ppm; see "Notes to RD" in memorandum: M. Sahafeyan, D35175, 5/7/08)

<u>Residues of Concern in Plants and Livestock</u>: Table 1 below contains a summary of tolerance expression and the residues of concern in plants and livestock commodities (for details see memorandums: G. Kramer, 22-JUL-1994; No DP# and W. Wassell, DP# 340379, 8/9/07).

Matrix	Residues of Concern			
	For Risk Assessment	For Tolerance Expression		
Plants	Parent Only	Parent Only		
Livestock	Parent and CGA 205375	Parent and CGA 205375		

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

Commodities		
Milk	Parent and CGA 205375	Parent and CGA 205375
Water	Parent Only	NA

<u>Tolerances</u>: HED recommends for the establishment of Section 18 tolerances on almond nutmeat at 0.05 ppm, almond hulls at 5.0 ppm and cucurbit vegetables (crop group 9) at 1.0 ppm in response to two Section 18 requests (discussed in memorandums with DP#s: D351715 and D351302); tolerances for livestock tissues are not changed by the addition of almond hulls to their diet (M. Sahafeyan, D351715, 7-MAY-08). The established and recommended tolerances are listed in Table 2 below.

Table 2. Tolerance Summary for Difenoconazole.						
Commodity	Established Tolerances (ppm)	Recommended Tolerances (ppm)				
Almond, nutmeat1	none	0.05				
Almond, hulls1	none	5.0				
Cucurbit Vegetables (Group 9)	none	1.0				
Fruit, Pome, (Group 11)	1.0					
Vegetable, Fruiting, Group 8	0.60					
Vegetable, Tuberous and Corm, subgroup 1C	0.01					
Beet, sugar <sup>2</sup>	0.30					
Papaya	0.30					
Apple, wet pomace	4.5					
Beet, sugar, dried pulp	1.9					
Potato, processed waste	0.04					
Banana	0.2					
Barley, grain	0.1					
Barley, hay	0.05					
Barley, straw	0.05					
Canola, seed	0.01					
Cattle fat	0.10					
Cattle, meat	0.05					
Cattle, meat byproducts	0.10					
Cattle, liver	0.20					
Corn, sweet, forage	0.01					
Corn, sweet, kernel plus cob with husks removed	0.01					
Corn, sweet, stover	0.01					
Cotton, gin byproducts	0.05					
Cotton, undelinted seed	0.05					
Egg	0.10					

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

	ary for Difenoconazole.	T
Commodity	Established	Recommended
~ ^	Tolerances (ppm)	Tolerances (ppm)
Goat, fat	0.10	
Goat, meat	0.05	
Goat, meat byproducts	0.10	
Goat, liver	0.20	
Grape	0.1	
Hog, fat	0.10	
Hog, meat	0.05	
Hog, meat byproducts	0.10	
Horse, fat	0.10	
Horse, meat	0.05	
Horse, meat byproducts	0.10	
Horse, liver	0.20	
Milk	0.01	
Rye, grain	0.1	
Sheep, fat	0.10	
Sheep, meat	0.05	
Sheep, meat byproducts	0.10	
Sheep, liver	0.20	
Wheat, forage	0.1	
Wheat, grain	0.1	
Wheat, straw	0.1	

Section 18 emergency exemption tolerance.

Food Residues and processing factors used in the Acute and Chronic Analysis: The acute and chronic analyses assumed tolerance-level residues for all commodities (crops and livestock tissues) and 100% CT for all the registered and proposed crops. Addition of almond hulls to the dietary burden of cattle did not result in an increase in the calculated maximum residues in beef and dairy cattle tissues. Experimental processing factors were used for apple juice (0.04x), potato chips (0.5x), potato granules/flakes (0.5x), sugar beet molasses (0.6x), sugar beet refined sugar (0.6x), tomato paste (1.6x), and tomato puree (0.5x); DEEM<sup>TM</sup> (ver. 7.81) default processing factors were assumed for other processed commodities.

## III. Drinking Water Data

The drinking water values used in the dietary risk assessment were provided by the Environmental Fate and Effects Division (EFED; Memo, I. Maher, 19-JUN-2007; DP# 333319). EFED conducted a Tier II drinking water assessment from surface water sources using Pesticide

<sup>&</sup>lt;sup>2</sup> The registered tolerance on sugar beet (0.01 ppm) needs to be changed to 0.3 ppm; see "Notes to RD" in memorandum: M. Sahafeyan, D351715, 07-MAY-08.

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

Root Zone/Exposure Analysis Modeling System (PRZM/EXAMS) model for the registered and proposed new uses and found that the highest estimated drinking water concentrations (EDWCs) are likely to be from surface water sources derived from aerial applications of difenoconazole to California ornamental nurseries at the maximum annual application rate of 0.53 lb ai/A. The estimated drinking water residues for 1-in-10 year annual peak, 1-in-10 year annual mean, and 36-year annual mean are 13.3, 9.43, and 7.18  $\mu$ g/L (ppb) respectively. The second highest EDWC was from Maine potatoes.

For ground water, SCI-GROW model indicates that difenoconazole concentrations are not likely to exceed 0.00128 µg/L (ppb) from either agricultural or non-agricultural uses.

In this assessment, 1-in-10-year annual peak (13.3 ppb), and 1-in-10-year annual mean (9.43 ppb) residue values were used for acute and chronic dietary exposure assessments respectively.

# IV. DEEM-FCID™ Program and Consumption Information

Difenoconazole acute and chronic dietary exposure assessments were conducted using the DEEM-FCID<sup>TM</sup> (ver. 2.03), which incorporates consumption data from USDA's CSFII (1994-1996 and 1998). The 1994-96, 98 data are based on the reported consumption of more than 20,000 individuals over two non-consecutive survey days. Foods "as consumed" (*i.e.*, apple pie) are linked to EPA-defined food commodities (*i.e.*, apples, peeled fruit - cooked; fresh or N/S; baked; or wheat flour - cooked; fresh or N/S, baked) using publicly available recipe translation files developed jointly by USDA/ARS and EPA. For chronic exposure assessment, consumption data are averaged for the entire U.S. population and within population subgroups, but for acute exposure assessment are retained as individual consumption events. Based on analysis of the 1994-96, 98 CSFII consumption data, which took into account dietary patterns and survey respondents, HED concluded that it is most appropriate to report risk for the following population subgroups: the general U.S. population, all infants (<1 year old), children 1-2, children 3-5, children 6-12, youth 13-19, adults 20-49, females 13-49, and adults 50+ years old.

For chronic dietary exposure assessment, an estimate of the residue level in each food or foodform (*i.e.*, orange or orange juice) on the food commodity residue list is multiplied by the average daily consumption estimate for that food/food form. The resulting residue consumption estimate for each food/food form is summed with the residue consumption estimates for all other food/food forms on the commodity residue list to arrive at the total average estimated exposure. Exposure is expressed in mg/kg body weight/day and as a percent of the cPAD. This procedure is performed for each population subgroup.

For acute exposure assessments, individual one-day food consumption data are used on an individual-by-individual basis. The reported consumption amounts of each food item can be multiplied by a residue point estimate and summed to obtain a total daily pesticide exposure for a deterministic exposure assessment, or "matched" in multiple random pairings with residue values and then summed in a probabilistic assessment. The resulting distribution of exposures is expressed as a percentage of the aPAD on both a user (*i.e.*, those who reported eating relevant commodities/food forms) and a per-capita (*i.e.*, those who reported eating the relevant

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

commodities as well as those who did not) basis. In accordance with HED policy, per capita exposure and risk are reported for all tiers of analysis. However, for Tiers 1 and 2, significant differences in user vs. per capita exposure and risk are identified and noted in the risk assessment.

## V. Toxicological Information

Table 3 below contains a summary of the relevant toxicological information for dietary risk analyses (M. Sahafeyan, D341303, 10/30/07)

Table 3. Summary of Toxicological Doses and Endpoints for Difenoconazole for Use in Dietary Risk Assessments.					
Exposure Scenario	Point of Departure	Uncertainty/FQPA Safety Factors	RfD, PAD for Risk Assessment	Study and Relevant Toxicological Effects	
Acute Dietary (All populations)	NOAEL = 25 mg/kg	$UF_{A} = 10X$ $UF_{H} = 10X$ $UF_{FQPA} = 1X$	aRfD = aPAD = 0.25 mg/kg/day	Acute Neurotoxicity Study in Rats  LOAEL= 200 mg/kg in males based on reduced fore-limb grip strength in males on day 1.	
Chronic Dietary (All populations)	NOAEL = 0.96 mg/kg/day	$UF_{A} = 10X$ $UF_{H} = 10X$ $UF_{FQPA} = 1X$	cRfD = cPAD = 0.01mg/kg/day	Combined chronic toxicity/carcinogenicity (rat; dietary) LOAEL = 24.1/32.8 mg/kg/day (M/F) based on cumulative decreases in body-weight gains.	
Cancer (oral, dermal, inhalation)  Difenoconazole is classified as a Group C, possible human carcinogen with a non-linear (MOE) approach for human risk characterization (CPRC Document, 7/27/94, Memo, P. V. Shah dated March 3, 2007, HED Doc. No. 0054532)					

Point of Departure (POD) = A data point or an estimated point that is derived from observed dose-response data and used to mark the beginning of extrapolation to determine risk associated with lower environmentally relevant human exposures. NOAEL = no observed adverse effect level. LOAEL = lowest observed adverse effect level. UF = uncertainty factor. UF<sub>A</sub> = extrapolation from animal to human (intraspecies). UF<sub>H</sub> = potential variation in sensitivity among members of the human population (interspecies). UF<sub>L</sub> = use of a LOAEL to extrapolate a NOAEL. UF<sub>S</sub> = use of a short-term study for long-term risk assessment. UF<sub>DB</sub> = to account for the absence of key data (i.e., lack of a critical study). FQPA SF = FQPA Safety Factor. PAD = population adjusted dose (a = acute, c = chronic). RfD = reference dose. MOE = margin of exposure.

#### VI. Results/Discussion

As stated above, for acute and chronic assessments, HED is concerned when dietary risk exceeds 100% of the aPAD or cPAD, respectively. The following paragraphs are summaries of the DEEM-FCID<sup>TM</sup> (ver. 2.03) acute and chronic exposure analyses.

Acute and chronic aggregate (food + water) analyses were performed using DEEM-FCID<sup>TM</sup> estimating the dietary exposure of the U.S. population and various population subgroups. The results are summarized in Tables 4 and 5 below for acute and chronic analyses respectively.

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

The resulting acute dietary (food + water) exposure estimates were less than HED's level of concern (<100% aPAD) at the 95<sup>th</sup> percentile of the exposure distribution for US general population (2.9% aPAD) and all population subgroups; the most highly-exposed population subgroup was all-infants <1 year old with 9.0% aPAD. The resulting chronic food exposure estimates were less than HED's level of concern (<100% cPAD) for U.S. general population (23% cPAD) and all population sub-groups; the most highly exposed population subgroup was children 1-2 years old with 65% cPAD. A cancer dietary assessment was not conducted for difenoconazole because the cancer NOAEL is higher than the chronic RfD; therefore, the chronic dietary risk estimate is more protective of any cancer effects.

Table 4. Summary of Dietary (Food and Drinking Water) Exposure and Risk for Difenoconazole						
	Acute Dietary (95th Percentile)		Chronic Dietary		Cancer	
Population Subgroup	Dietary Exposure (mg/kg/day)	% aPAD	Dietary Exposure (mg/kg/day)	% cPAD	Dietary Exposure (mg/kg/day)	Risk
General U.S. Population	0.007277	2.9	0.002265	23		
All Infants (< 1 year old)	0.022518	9.0	0.005586	56		
Children 1-2 years old	0.019060	7.6	0.006516	65	·	
Children 3-5 years old	0.016481	6.6	0.005547	56		
Children 6-12 years old	0.009954	4.0	0.003287	33		27/4
Youth 13-19 years old	0.004989	2.0	0.001663	17	N/A	N/A
Adults 20-49 years old	0.004768	1.9	0.001632	16	·	
Adults 50+ years old	0.005685	2.3	0.001880	19		
Females 13-49 years old	0.004915	2.0	0.001637	16		

The bolded %cPAD is the highest.

### VII. Characterization of Inputs/Outputs

The acute and chronic analyses assumed tolerance-level residues, 100% CT, and empirical and DEEM™ default processing factors. Therefore, these analyses were considered conservative and could be further refined through the use of anticipated residues (ARs) for all commodities, % market share data for the proposed commodities, %CT data for registered commodities, and/or empirical processing factors for all commodities.

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

#### VIII. Conclusions

Acute and chronic aggregate (food + water) dietary risk assessments were conducted for difenoconazole using the DEEM-FCID<sup>TM</sup> (ver. 2.03) model and assumed tolerance-level residues, 100% CT, and empirical and DEEM<sup>TM</sup> default processing factors. The resulting acute and chronic aggregate exposure estimates were less than HED's level of concern, and therefore are not of concern. The most highly-exposed population subgroups in the acute (at the 95<sup>th</sup> percentile of the exposure distribution) and chronic analyses were all-infants <1 year old (9.0% aPAD) and children 1-2 years old (65% cPAD) respectively.

#### IX. Attachments

Attachment 1: DEEM-FCID™ Acute Residue File

Attachment 2: DEEM-FCID™ Acute Exposure Estimates
Attachment 3: DEEM-FCID™ Chronic Residue File

Attachment 4: DEEM-FCID™ Chronic Exposure Estimates

RDI: DESAC (23-APR-2008)

Registration Number(s): 08CA12 and 08GA01

DP#: 351961 and 352204

PC Code: 128847

M. Sahafeyan: S10944:PY1:(703)-305-0776

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

## Attachment 1: DEEM-FCID™ Acute Residue File

Filename: C:\Documents and Settings\msahafey\My Documents\difenconazole\Final

Documents\Section 18- almonds\difenoconazole\_acute-Sec18-043008.R98

Chemical: Difenoconazole

RfD(Chronic): .01 mg/kg bw/day NOEL(Chronic): 0 mg/kg bw/day RfD(Acute): .25 mg/kg bw/day NOEL(Acute): 0 mg/kg bw/day Date created/last modified: 04-29-2008/10:54:48/8

Program ver. 2.03

		last modified: 04-29-2008/10:54:4			
EPA Comment	Crop		Def Res	Adj.Fa	ctors
Code	Grp	Commodity Name	(mgg)	#1	#2
L4000030	14	Almond	0.050000	1.000	1.000
4000031		Almond-babyfood	0.050000	1.000	1.000
4000040		Almond. oil	0.050000	1.000	1.000
14000041		Almond, oil-babyfood	0.050000	1.000	1.000
1000070		Apple, fruit with peel	1.000000	1.000	1.000
1000080		Apple, peeled fruit	1.000000	1.000	1.000
1000081		Apple, peeled fruit Apple, peeled fruit-babyfood	1.000000	1.000	1.000
1000090		Apple, dried	1.000000	8.000	1.000
1000091		Apple, dried-babyfood	1.000000	8.000	1.000
1000001		Apple, juice	1.000000	0.040	1.000
1000100		Apple, juice	1.000000	0.040	1.000
1000101		Apple, juice-babyfood Apple, sauce	1.000000	1.000	1.000
11000110		Apple, sauce-babyfood	1.000000	1.000	1.000
			1.000000	1.000	1.000
09020210		Balsam pear			
95000230		Banana	0.200000	1.000	1.000
95000231		Banana-babyfood	0.200000	1.000	1.000
95000240		Banana, dried	0.200000	3.900	1.000
95000241		Banana, dried-babyfood	0.200000	3.900	1.000
L5000250		Barley, pearled barley	0.100000	1.000	1.000
L5000251		Barley, pearled barley-babyfood	0.100000	1.000	1.000
L5000260		Barley, flour	0.100000	1.000	1.000
15000261		Barley, flour-babyfood	0.100000	1.000	1.000
L5000270		Barley, bran	0.100000	1.000	1.000
21000440		Beef, meat	0.050000	1.000	1.000
21000441		Beef, meat-babyfood Beef, meat, dried Beef meat, byproducts	0.050000	1.000	1.000
21000450		Beef, meat, dried	0.050000	1.920	1.000
21000460		Deel, meat Dybloducts	0.100000	1.000	1.000
21000461		Beef, meat byproducts-babyfood	0.100000	1.000	1.000
21000470		Beef, fat	0.100000	1.000	1.000
21000471	M	Beef,fat-babyfood	0.100000	1.000	1.000
21000480	M	Beef, kidney	0.100000	1.000	1.000
21000490	M	Beef, liver	0.200000	1.000	1.000
21000491		Beef, liver-babyfood	0.200000		1.000
01010520	1A	Beet, sugar	0.300000	1.000	1.000
01010521	1A	Beet, sugar-babyfood	0.300000	1.000	1.000
01010530	1A	Beet, sugar, molasses	0.300000	0.600	1.000
01010531	1A	Beet, sugar, molasses-babyfood	0.300000	0.600	1.000
9010750		Cantaloupe	1.000000	1.000	1.000
9010800		Casaba	1.000000	1.000	1.000
9020880		Chayote, fruit	1.000000	1.000	1.000
09021020		Chinese waxgourd	1.000000	1.000	1.000
15001270		Corn, sweet	0.010000	1.000	1.000
15001270		Corn, sweet-babyfood	0.010000	1.000	1.000
95001271		Cottonseed, oil	0.050000	1.000	1.000
95001280		Cottonseed, oil-babyfood	0.050000	1.000	1.000
1001290		Crabapple	1.000000	1.000	1.000
1 1 (1) 1 7 9 11					

Difenoconazole PC Code: 128847	Dietary Exposure and Risk Assessment		DP#: 35	1961 and 35220
70001450 P	Egg, whole	0.100000	1.000	1.000
70001451 P	Egg, whole-babyfood	0.100000	1.000	1.000
70001460 P	Egg, white	0.100000	1.000	1.000
70001461 P	Egg, white (solids)-babyfood	0.100000	1.000	1.000
70001470 P	Egg, yolk	0.100000	1.000	1.000
70001471 P	Egg, yolk-babyfood	0.100000 0.050000	1.000 1.000	1.000
23001690 M 23001700 M	Goat, meat Goat, meat byproducts	0.100000	1.000	1.000
23001700 M 23001710 M	Goat, fat	0.100000	1.000	1.000
23001710 M 23001720 M	Goat, kidney	0.100000	1.000	1.000
23001730 M	Goat, liver	0.200000	1.000	1.000
95001750 O	Grape	0.100000	1.000	1.000
95001760 O	Grape, juice	0.100000	1.200	1.000
95001761 O	Grape, juice-babyfood	0.100000	1.200	1.000
95001770 0	Grape, leaves	0.100000	1.000	1.000
95001780 0	Grape, raisin	0.100000 0.100000	4.300 1.000	1.000 1.000
95001790 O 09011870 9A	Grape, wine and sherry Honeydew melon	1.000000	1.000	1.000
24001890 M	Horse, meat	0.050000	1.000	1.000
11002100 11	Loquat	1.000000	1.000	1.000
27002220 D	Milk, fat	0.010000	1.000	1.000
27002221 D	Milk, fat - baby food/infant for	0.010000	1.000	1.000
27012230 D	Milk, nonfat solids	0.010000	1.000	1.000
27012231 D	Milk, nonfat solids-baby food/in	0.010000	1.000	1.000
27022240 D	Milk, water	0.010000	1.000	1.000
27022241 D	Milk, water-babyfood/infant form	0.010000	1.000	1.000
27032251 D	Milk, sugar (lactose)-baby food/ Okra	0.010000 0.600000	1.000	1.000 1.000
08002340 8 95002450 O	Papaya	0.300000	1.000	1.000
95002451 0	Papaya-babyfood	0.300000	1.000	1.000
95002460 O	Papaya, dried	0.300000	1.800	1.000
95002470 O	Papaya, juice	0.300000	1.500	1.000
11002660 11	Pear	1.000000	1.000	1.000
11002661 11	Pear-babyfood	1.000000	1.000	1.000
11002670 11	Pear, dried	1.000000 1.000000	6.250	1.000
11002680 11	Pear, juice Pear, juice-babyfood	1.000000	$1.000 \\ 1.000$	1.000 1.000
11002681 11 08002700 8	Pepper, bell	0.600000	1.000	1.000
08002701 8	Pepper, bell-babyfood	0.600000		
08002710 8	Pepper, bell, dried	0.600000	1.000	1.000
08002711 8	Pepper, bell, dried-babyfood	0.600000	1.000	1.000
08002720 8	Pepper, nonbell	0.600000	1.000	1.000
08002721 8	Pepper, nonbell-babyfood	0.600000	1.000	1.000
08002730 8	Pepper, nonbell, dried	0.600000 0.200000	1.000	1.000 1.000
95002830 0	Plantain Plantain, dried	0.200000	3.900	1.000
95002840 O 25002900 M	Pork, meat	0.050000	1.000	1.000
25002900 M	Pork, meat-babyfood	0.050000	1.000	1.000
25002910 M	Pork, skin	0.100000	1.000	1.000
25002920 M	Pork, meat byproducts	0.100000	1.000	1.000
25002921 M	Pork, meat byproducts-babyfood	0.100000	1.000	1.000
25002930 M	Pork, fat	0.100000	1.000	1.000
25002931 M	Pork, fat-babyfood	0.100000	1.000	1.000
25002940 M	Pork, kidney	0.100000 0.200000	1.000 1.000	1.000 1.000
25002950 M 01032960 1C	Pork, liver Potato, chips	0.010000	0.500	1.000
01032960 1C 01032970 1C	Potato, dry (granules/ flakes)	0.010000	0.500	1.000
01032970 1C 01032971 1C	Potato, dry (granules/ flakes)-b	0.010000	0.500	1.000
01032972 1C	Potato, flour	0.010000	1.000	1.000
01032981 1C	Potato, flour-babyfood	0.010000	1.000	1.000
01032990 1C	Potato, tuber, w/peel	0.010000	1.000	1.000
01032991 1C	Potato, tuber, w/peel-babyfood	0.010000	1.000	1.000

Difenoconazole PC Code: 128847	Dietary Exposure and Risk Assessment			1961 and 352204
01033000 1C	Potato, tuber, w/o peel	0.010000	1.000	1.000
01033000 1C 01033001 1C	Potato, tuber, w/o peel-babyfood	0.010000	1.000	1.000
09023080 9B	Pumpkin	1.000000	1.000	1.000
09023090 9B	Pumpkin, seed	1.000000	1.000	1.000
11003100 11	Ouince	1.000000	1.000	1.000
20003190 20	Rapeseed, oil	0.010000	1.000	1.000
20003191 20	Rapeseed, oil-babyfood	0.010000	1.000	1.000
15003280 15	Rye, grain	0.100000	1.000	1.000
15003290 15	Rye, flour	0.100000	1.000	1.000
26003390 M	Sheep, meat	0.050000	1.000	1.000
26003391 M	Sheep, meat-babyfood	0.050000	1.000	1.000
26003400 M	Sheep, meat byproducts	0.100000	1.000	1.000
26003410 M	Sheep, fat	0.100000	1.000	1.000
26003411 M	Sheep, fat-babyfood	0.100000	1.000	1.000
26003420 M	Sheep, kidney	0.100000	1.000	1.000
26003430 M	Sheep, liver	0.200000	1.000	1.000
09023560 9B	Squash, summer	1.000000	1.000	1.000
09023561 9B	Squash, summer-babyfood	1.000000	1.000	1.000
09023570 9B	Squash, winter	1.000000	1.000	1.000
09023571 9B	Squash, winter-babyfood	1.000000	1.000	1.000
08003740 8	Tomatillo	0.600000	1.000	1.000
08003750 8	Tomato	0.600000	1.000	1.000
08003751 8	Tomato-babyfood	0.600000	1.000	1.000
08003760 8	Tomato, paste	0.600000	1.600	1.000
08003761 8	Tomato, paste-babyfood	0.600000	1.600	1.000
08003770 8	Tomato, puree	0.600000	0.500	1.000
08003771 8	Tomato, puree-babyfood	0.600000	0.500	1.000
08003780 8	Tomato, dried	0.600000	14.300	1.000
08003781 8	Tomato, dried-babyfood	0.600000	14.300	1.000
08003790 8	Tomato, juice	0.600000	1.500	1.000
86010000 0	Water, direct, all sources	0.013300	1.000	1.000
86020000 O	Water, indirect, all sources	0.013300	1.000 1.000	1.000 1.000
09013990 9A	Watermelon	1.000000	1.000	1.000
09014000 9A 15004010 15	Watermelon, juice Wheat, grain	0.100000	1.000	1.000
	Wheat, grain-babyfood	0.100000	1.000	1.000
15004011 15 15004020 15	Wheat, flour	0.100000		1.000
15004020 15	Wheat, flour-babyfood	0.100000	1.000	1.000
15004021 15	Wheat, germ	0.100000	1.000	1.000
15004030 15	Wheat, bran	0.100000	1.000	1.000
15004040 15	wneat, bran	0.100000	1.000	1.000

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

## Attachment 2: DEEM-FCID™ Acute Exposure Estimates

U.S. Environmental Protection Agency

Ver. 2.02

DEEM-FCID ACUTE Analysis for DIFENOCONAZOLE

(1994-98 data)

Residue file: difenoconazole\_acute-Sec18-043008.R98

Adjustment factor #2 NOT used.

Residue file dated: 04-30-2008/11:37:24/8

Analysis Date: 04-30-2008/11:44:37 Residue file do Daily totals for food and foodform consumption used.

Run Comment: "Two section 18 requests were included: almond nutmeat (0.05 ppm)

and cucurbit crop group 9 (1.0 ppm); livestock are not affected."

#### Summary calculations (per capita):

	95th Perc Exposure	entile % aRfD	99th Perc Exposure		99.9th Pe Exposure	rcentile % aRfD
U.S. Population:	0.007277	2.91	0.015755	6.30	0.035323	14.13
All infants:	0.022518	9.01	0.035271	14.11	0.060480	24.19
Children 1-2 yrs:	0.019060	7.62	0.031084	12.43	0.063303	25.32
Children 3-5 yrs:	0.016481	6.59	0.029090	11.64	0.055217	22.09
Children 6-12 yrs:	0.009954	3.98	0.019901	7.96	0.043000	17.20
Youth 13-19 yrs:	0.004989	2.00	0.010215	4.09	0.027368	10.95
Adults 20-49 yrs:	0.004768	1.91	0.008419	3.37	0.018473	7.39
Adults 50+ yrs:	0.005685	2.27	0.009946	3.98	0.022200	8.88
Females 13-49 yrs:	0.004915	1.97	0.008969	3.59	0.020682	8.27

Dietary Exposure and Risk Assessment

PC Code: 128847

DP#: 351961 and 352204

## Attachment 3: DEEM-FCID™ Chronic Residue File

Filename: C:\Documents and Settings\msahafey\My Documents\difenconazole\Final

Documents\Section 18- almonds\difenoconazole\_chronic-Sec18-050608.R98

Chemical: Difenoconazole

Difenoconazole

 $\label{eq:reduced_reduced_reduced_reduced} $$RfD(Chronic): 0.01 \ mg/kg \ bw/day \ NOEL(Chronic): 4.7 \ mg/kg \ bw/day \ RfD(Acute): .25 \ mg/kg \ bw/day \ NOEL(Acute): .25$ 

Date created/last modified: 05-06-2008/12:34:16/8 Program ver. 2.03

EPA Comment	Crop		Def Res	Adj.Fa	ctors
Code	Grp	Commodity Name	(ppm)	#1	#2
14000030	14	Almond	0.050000	1.000	1.000
14000031	14	Almond-babyfood	0.050000	1.000	1.000
14000040	14	Almond, oil	0.050000	1.000	1.000
14000041	14	Almond, oil-babyfood	0.050000	1.000	1.000
11000070		Apple, fruit with peel	1.000000	1.000	1.000
11000080	11	Apple, peeled fruit	1.000000	1.000	1.000
11000081		Apple, peeled fruit-babyfood	1.000000	1.000	1.000
11000090	11	Apple, dried	1.000000	8.000	1.000
11000091	11	Apple, dried-babyfood	1.000000	8.000	1.000
11000100	11	Apple, juice	1.000000	0.040	1.000
11000101	11	Apple, juice-babyfood	1.000000	0.040	1.000
11000110	11	Apple, sauce	1.000000	1.000	1.000
11000111	11	Apple, sauce-babyfood	1.000000	1.000	1.000
09020210	9B	Balsam pear	1.000000	1.000	1.000
95000230	0	Banana	0.200000	1.000	1.000
95000231	0	Banana-babyfood	0.200000	1.000	1.000
95000240	0	Banana, dried	0.200000	3.900	1.000
95000241	0	Banana, dried-babyfood	0.200000	3.900	1.000
15000250	15	Barley, pearled barley	0.100000	1.000	1.000
15000251	15	Barley, pearled barley-babyfood	0.100000	1.000	1.000
15000260	15	Barley, flour	0.100000	1.000	1.000
15000261	15	Barley, flour-babyfood	0.100000	1.000	1.000
15000270	15	Barley, bran	0.100000	1.000	1.000
21000440	M	Beef, meat	0.050000	1.000	1.000
21000441	M	Beef, meat-babyfood	0.050000	1.000	1.000
21000450	M	Beef, meat, dried	0.050000	1.920	1.000
21000460	M	Beef, meat byproducts	0.100000	1.000	1.000
21000461	M	Beef, meat byproducts-babyfood	0.100000	1.000	1.000
21,000470	M	Beef, fat	0.100000	1.000	1.000
21000471	M	Beef,fat-babyfood	0.100000	1.000	1.000
21000480	M	Beef, kidney	0.100000	1.000	1.000
21000490	M	Beef, liver	0.200000	1.000	1.000
21000491	M	Beef, liver-babyfood	0.200000	1.000	1.000
01010520	1A	Beet, sugar	0.300000	1.000	1.000
01010521	1A	Beet, sugar-babyfood	0.300000	1.000	1.000
01010530		Beet, sugar, molasses	0.300000	0.600	1.000
01010531		Beet, sugar, molasses-babyfood	0.300000	0.600	1.000
09010750	9A	Cantaloupe	1.000000	1.000	1.000
09010800		Casaba	1.000000	1.000	1.000
09020880		Chayote, fruit	1.000000	1.000	1.000
09021020		Chinese waxgourd	1.000000	1.000	1.000
15001270		Corn, sweet	0.010000	1.000	1.000
15001271		Corn, sweet-babyfood	0.010000	1.000	1.000
95001280		Cottonseed, oil	0.050000	1.000	1.000
95001281		Cottonseed, oil-babyfood	0.050000	1.000	1.000

Difenoconazole PC Code: 128847	Dietary Exposure and Risk Assessmen	t .	DP#: 351961 and 352204
11001290 11	Crabapple	1.000000	1.000 1.000
09021350 9B	Cucumber	1.000000	1.000 1.000
70001450 P	Egg, whole	0.100000	1.000 1.000
70001451 P	Egg, whole-babyfood	0.100000	1.000 1.000
70001460 P	Egg, white	0.100000	1.000 1.000
70001461 P	Egg, white (solids)-babyfood	0.100000	1.000 1.000
70001470 P	Egg, yolk	0.100000	1.000 1.000
70001471 P	Egg, yolk-babyfood	0.100000	1.000 1.000
23001690 M	Goat, meat	0.050000	1.000 1.000
23001700 M	Goat, meat byproducts	0.100000	1.000 1.000
23001710 M	Goat, fat	0.100000	1.000 1.000
23001720 M 23001730 M	Goat, kidney Goat, liver	0.100000 0.200000	1.000 1.000 1.000 1.000
95001750 M	Grape	0.100000	1.000 1.000
95001750 O	Grape, juice	0.100000	1.200 1.000
95001761 O	Grape, juice-babyfood	0.100000	1.200 1.000
95001770 O	Grape, leaves	0.100000	1.000 1.000
95001780 O	Grape, raisin	0.100000	4.300 1.000
95001790 O	Grape, wine and sherry	0.100000	1.000 1.000
09011870 9A	Honeydew melon	1.000000	1.000 1.000
24001890 M	Horse, meat	0.050000	1.000 1.000
11002100 11	Loquat	1.000000	1.000 1.000
27002220 D	Milk, fat	0.010000	1.000 1.000
27002221 D	Milk, fat - baby food/infant for	0.010000	1.000 1.000
27012230 D	Milk, nonfat solids	0.010000	1.000 1.000
27012231 D	Milk, nonfat solids-baby food/in	0.010000	1.000 1.000
27022240 D	Milk, water	0.010000	1.000 1.000
27022241 D	Milk, water-babyfood/infant form	0.010000	1.000 1.000
27032251 D	Milk, sugar (lactose)-baby food/	0.010000	1.000 1.000
08002340 8	Okra	0.600000	1.000 1.000
95002450 0	Papaya	0.300000	1.000 1.000
95002451 O 95002460 O	Papaya-babyfood Papaya, dried	0.300000 0.300000	1.000 1.000 1.800 1.000
95002470 O	Papaya, juice	0.300000	1.500 1.000
11002660 11	Pear	1.000000	1.000 1.000
11002661 11	Pear-babyfood	1.000000	1.000 1.000
11002670 11	Pear, dried	1.000000	6.250 1.000
11002680 11	Pear, juice	1.000000	1.000 1.000
11002681 11	Pear, juice-babyfood	1.000000	1.000 1.000
08002700 8	Pepper, bell	0.600000	1.000 1.000
08002701 8	Pepper, bell-babyfood	0.600000	1.000 1.000
08002710 8	Pepper, bell, dried	0.600000	1.000 1.000
08002711 8	Pepper, bell, dried-babyfood	0.600000	1.000 1.000
08002720 8	Pepper, nonbell	0.600000	1.000 1.000
08002721 8	Pepper, nonbell-babyfood	0.600000	1.000 1.000
08002730 8	Pepper, nonbell, dried	0.600000	1.000 1.000
95002830 O	Plantain Plantain, dried	0.200000 0.200000	1.000 1.000
95002840 O 25002900 M	Pork, meat	0.050000	3.900 1.000 1.000 1.000
25002900 M 25002901 M	Pork, meat-babyfood	0.050000	1.000 1.000
25002910 M	Pork, skin	0.100000	1.000 1.000
25002920 M	Pork, meat byproducts	0.100000	1.000 1.000
25002921 M	Pork, meat byproducts-babyfood	0.100000	1.000 1.000
25002930 M	Pork, fat	0.100000	1.000 1.000
25002931 M	Pork, fat-babyfood	0.100000	1.000 1.000
25002940 M	Pork, kidney	0.100000	1.000 1.000
25002950 M	Pork, liver	0.200000	1.000 1.000
01032960 1C	Potato, chips	0.010000	0.500 1.000
01032970 1C	Potato, dry (granules/ flakes)	0.010000	0.500 1.000
01032971 1C	Potato, dry (granules/ flakes)-b	0.010000	0.500 1.000
01032980 1C	Potato, flour	0.010000	1.000 1.000
01032981 1C	Potato, flour-babyfood	0.010000	1.000 1.000

Difenoconazole PC Code: 128847	Dietary Exposure and Risk Assessment		DP#: 351961 and 352204	
01032990 1C	Potato, tuber, w/peel	0.010000	1.000	1.000
01032991 1C	Potato, tuber, w/peel-babyfood	0.010000	1.000	1.000
01032991 1C	Potato, tuber, w/o peel	0.010000	1.000	1.000
01033000 1C	Potato, tuber, w/o peel-babyfood	0.010000	1.000	1.000
09023080 9B	Pumpkin	1.000000	1.000	1.000
09023090 9B	Pumpkin, seed	1.000000	1.000	1.000
11003100 11	Quince	1.000000	1.000	1.000
20003190 20	Rapeseed, oil	0.010000	1.000	1.000
20003190 20	Rapeseed, oil-babyfood	0.010000	1.000	1.000
15003280 15	Rye, grain	0.100000	1.000	1.000
15003290 15	Rye, flour	0.100000	1.000	1.000
26003390 M	Sheep, meat	0.050000	1.000	1.000
26003391 M	Sheep, meat-babyfood	0.050000	1.000	1.000
26003400 M	Sheep, meat byproducts	0.100000	1.000	1.000
26003410 M	Sheep, fat	0.100000	1.000	1.000
26003411 M	Sheep, fat-babyfood	0.100000	1.000	1.000
26003420 M	Sheep, kidney	0.100000	1.000	1.000
26003430 M	Sheep, liver	0.200000	1.000	1.000
09023560 9B	Squash summer	1.000000	1.000	1.000
09023561 9B	Squash, summer-babyfood	1.000000	1.000	1.000
09023570 9B	Squash, winter	1.000000	1.000	1.000
09023571 9B	Squash, winter-babyfood		1.000	1.000
08003740 8	Tomatillo	0.600000	1.000	1.000
08003750 8	Tomato	0.600000	1.000	1.000
08003751 8	Tomato-babyfood	0.600000	1.000	1.000
08003760 8	Tomato, paste	0.600000	1.600	1.000
08003761 8	Tomato, paste-babyfood	0.600000	1.600	1.000
08003770 8	Tomato, puree	0.600000	0.500	1.000
08003771 8	Tomato, puree-babyfood	0.600000	0.500	1.000
08003780 8	Tomato, dried	0.600000	14.300	1.000
08003781 8	Tomato, dried-babyfood	0.600000	14.300	1.000
08003790 8	Tomato, juice	0.600000	1.500	1.000
86010000 O	Water, direct, all sources	0.009430	1.000	1.000
86020000 O	Water, indirect, all sources	0.009430	1.000	1.000
09013990 9A	Watermelon	1.000000	1.000	1.000
09014000 9A	Watermelon, juice	1.000000	1.000	1.000
15004010 15	Wheat, grain	0.100000	1.000	1.000
15004011 15	Wheat, grain-babyfood	0.100000	1.000	1.000
15004020 15	Wheat, flour	0.100000	1.000	1.000
15004021 15	Wheat, flour-babyfood	0.100000	1.000	1.000
15004030 15	Wheat, germ	0.100000	1.000	1.000
15004040 15	Wheat, bran	0.100000	1.000	1.000

Dietary Exposure and Risk Assessment

DP#: 351961 and 352204

# Attachment 4: DEEM-FCID™ Chronic Exposure Estimates

U.S. Environmental Protection Agency

Ver. 2.00

DEEM-FCID Chronic analysis for DIFENOCONAZOLE

(1994-98 data)

Residue file name: C:\Documents and Settings\msahafey\My Documents\difenconazole\Final

Documents\Section 18- almonds\difenoconazole\_chronic-Sec18-050608.R98

Adjustment factor #2 NOT used.

Analysis Date 05-06-2008/12:35:21

Residue file dated: 05-06-2008/12:34:16/8

Reference dose (RfD, Chronic) = .01 mg/kg bw/day

Total exposure by population subgroup

#### Total Exposure \_\_\_\_\_

Population Subgroup	mg/kg body wt/day	Percent of Rfd	
U.S. Population (total)	0.002265	22.6%	
U.S. Population (spring season) U.S. Population (summer season) U.S. Population (autumn season) U.S. Population (winter season)	0.002282 0.002534 0.002201 0.002030	22.8% 25.3% 22.0% 20.3%	
Northeast region Midwest region Southern region Western region	0.002352 0.002285 0.001996 0.002593	23.5% 22.8% 20.0% 25.9%	
Hispanics Non-hispanic whites Non-hispanic blacks Non-hisp/non-white/non-black	0.002606 0.002232 0.001929 0.002949	26.1% 22.3% 19.3% 29.5%	
All infants (< 1 year) Nursing infants Non-nursing infants Children 1-6 yrs Children 7-12 yrs	0.005586 0.003486 0.006383 0.005675 0.003091	55.9% 34.9% 63.8% 56.8% 30.9%	
Females 13-19 (not preg or nursing) Females 20+ (not preg or nursing) Females 13-50 yrs Females 13+ (preg/not nursing) Females 13+ (nursing)	0.001532 0.001746 0.001819 0.002087 0.002461	15.3% 17.5% 18.2% 20.9% 24.6%	
Males 13-19 yrs Males 20+ yrs Seniors 55+	0.001788 0.001680 0.001925	17.9% 16.8% 19.3%	
Children 1-2 yrs Children 3-5 yrs Children 6-12 yrs Youth 13-19 yrs Adults 20-49 yrs Adults 50+ yrs Females 13-49 yrs	0.006516 0.005547 0.003287 0.001663 0.001632 0.001880 0.001637	65.2% 55.5% 32.9% 16.6% 16.3% 18.8%	



# R159255

Chemical Name: Difenoconazole

PC Code: 128847

**HED File Code:** 11000 Chemistry Reviews

Memo Date: 5/1/2008

File ID: DPD351961

DPD352204 DPD322238 DPD351751 DPD341303 DPD340379 DPD333319

Accession #: 000-00-0125

**HED Records Reference Center** 

7/2/2008